



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Sex-specific clines support incipient speciation in a common European mammal

Sutter, Andreas ; Beysard, Mathias ; Heckel, Gerald

Abstract: Hybrid zones provide excellent opportunities to study processes and mechanisms underlying reproductive isolation and speciation. Here we investigated sex-specific clines of molecular markers in hybrid zones of morphologically cryptic yet genetically highly-diverged evolutionary lineages of the European common vole (*Microtus arvalis*). We analyzed the position and width of four secondary contact zones along three independent transects in the region of the Alps using maternally (mitochondrial DNA) and paternally (Y-chromosome) inherited genetic markers. Given male-biased dispersal in the common vole, a selectively neutral secondary contact would show broader paternal marker clines than maternal ones. In a selective case, for example, involving a form of Haldane's rule, Y-chromosomal clines would not be expected to be broader than maternal markers because they are transmitted by the heterogametic sex and thus gene flow would be restricted. Consistent with the selective case, paternal clines were significantly narrower or at most equal in width to maternal clines in all contact zones. In addition, analyses using maximum likelihood cline-fitting detected a shift of paternal relative to maternal clines in three of four contact zones. These patterns suggest that processes at the contact zones in the common vole are not selectively neutral, and that partial reproductive isolation is already established between these evolutionary lineages. We conclude that hybrid zone movement, sexual selection and/or genetic incompatibilities are likely associated with an unusual unidirectional manifestation of Haldane's rule in this common European mammal.

DOI: <https://doi.org/10.1038/hdy.2012.124>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-77021>

Journal Article

Accepted Version

Originally published at:

Sutter, Andreas; Beysard, Mathias; Heckel, Gerald (2013). Sex-specific clines support incipient speciation in a common European mammal. *Heredity*, 110(4):398-404.

DOI: <https://doi.org/10.1038/hdy.2012.124>

Sex-specific clines support incipient speciation in a common European mammal

Andreas Sutter¹⁾, Mathias Beysard^{1), 2)} and Gerald Heckel^{1), 2)}

¹⁾ Computational and Molecular Population Genetics (CMPG), Institute of Ecology and Evolution, University of Bern, CH 3012 Bern, Switzerland

²⁾ Swiss Institute of Bioinformatics, Genopode, CH 1015 Lausanne, Switzerland

Correspondence:

Gerald Heckel, Computational and Molecular Population Genetics (CMPG), Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, CH 3012 Bern, Switzerland
phone: +41 31 631 30 29, fax: +41 31 631 48 88, email: gerald.heckel@iee.unibe.ch

Running title: Incipient speciation in the common vole

Keywords: cline analyses, *Microtus arvalis*, secondary contact zone, Haldane's rule, Y-chromosome, speciation

Word count: 5863

23 **Abstract**

24 Hybrid zones provide excellent opportunities to study processes and mechanisms
25 underlying reproductive isolation and speciation. Here we investigated sex-specific clines of
26 molecular markers in hybrid zones of morphologically cryptic yet genetically highly-diverged
27 evolutionary lineages of the European common vole (*Microtus arvalis*). We analyzed the
28 position and width of four secondary contact zones along three independent transects in the
29 region of the Alps using maternally (mitochondrial DNA) and paternally (Y chromosome)
30 inherited genetic markers. Given male-biased dispersal in the common vole, a selectively
31 neutral secondary contact would show broader paternal marker clines than maternal ones. In a
32 selective case e.g. involving a form of Haldane's rule, Y-chromosome clines would not be
33 expected to be broader than maternal markers because they are transmitted by the
34 heterogametic sex and thus gene flow would be restricted. Consistent with the selective case,
35 paternal clines were significantly narrower or at most equal in width to maternal clines in all
36 contact zones. Additionally, analyses using maximum likelihood cline-fitting detected a shift of
37 paternal relative to maternal clines in three of four contact zones. These patterns suggest that
38 processes at the contact zones in the common vole are not selectively neutral and that partial
39 reproductive isolation is already established between these evolutionary lineages. We conclude
40 that hybrid zone movement, sexual selection and genetic incompatibilities are likely associated
41 with an unusual unidirectional manifestation of Haldane's rule in this common European
42 mammal.

43 **Introduction**

44 Speciation involves the establishment of reproductive isolation between organisms.
45 Reproductive isolation may be achieved through various mechanisms acting singly or together
46 at the pre- and/or post-zygotic stage, such as assortative mating or genetic incompatibilities
47 (Coyne and Orr, 2004; Turelli and Begun, 1997). Incomplete reproductive isolation is often
48 characterized by sex-specific processes, and in particular intrinsic genetic incompatibilities often
49 more strongly affect hybrids of the heterogametic sex, a pattern known as Haldane's rule
50 (Haldane, 1922; Schilthuizen *et al.*, 2011). One expectation from Haldane's rule is that the
51 extent of gene flow between incipient species may strongly differ if a gene is paternally or
52 maternally transmitted (Coyne and Orr, 2004; Macholan *et al.*, 2007; Turelli and Begun, 1997).

53 Hybrid zones offer the possibility to study the extent and potential sex-specificity of
54 gene flow between incipient species in their natural environment. Here the consistency of
55 introgression of genetic markers may provide valuable information on the degree of
56 reproductive isolation between organisms (Barton and Hewitt, 1989; Raufaste *et al.*, 2005;
57 Schilthuizen *et al.*, 2011). For example, in a hybrid zone with a high degree of reproductive
58 isolation, Haldane's rule might lead to very low gene flow for male-specific Y-chromosomal
59 markers in mammals whereas no impediments for gene flow might exist for maternally
60 transmitted mitochondrial markers (e.g. Beysard *et al.*, 2012). Despite the recognition of the
61 crucial information carried by the Y chromosome in mammals (Petit *et al.*, 2002), its
62 investigation in hybrid zones of non-model organisms remains restricted to very few systems

(e.g. Hellborg *et al.*, 2005; Jaarola *et al.*, 1997; Yannic *et al.*, 2008), most probably because of technical limitations.

The rich technical and genomic resources available for the laboratory mouse have eased the development of the secondary contact zone between the house mouse sub-species *Mus musculus musculus* and *Mus musculus domesticus* running through Europe to a potential model system for speciation research in mammals. The exploration of this contact zone has shown e.g. that the subspecies may mate assortatively (see Ganem *et al.*, 2008), hybrids may differ from parentals in parasite load (see Baird *et al.*, 2012) or male hybrids may be sterile (reviewed in Britton-Davidian *et al.*, 2005; see also Mihola *et al.*, 2009). The observation of male sterility combined with the detection of abrupt Y and X chromosome marker clines (Dod *et al.*, 1993; Macholan *et al.*, 2007; 2008; 2011) relative to other markers across the hybrid zone are consistent with the hypothesis that sex chromosomes and in particular factors on the X chromosome play an important role in causing genetic incompatibilities in mouse hybrids (Coyne and Orr, 2004). Nevertheless, there is extensive geographical and individual variation in male sterility (Britton-Davidian *et al.*, 2005; Good *et al.*, 2008; Turner *et al.*, 2012), and sex-chromosomal marker clines may be not coincident with other genetic markers in particular geographic regions, potentially due to asymmetrical processes (Jones *et al.*, 2010; Macholan *et al.*, 2008). For example, the Y-chromosome was found non-coincident with other markers in an area of about 330 km² in the Czech-Bavarian part of the mouse hybrid zone (Macholan *et al.*, 2008). This may be explained by two non-mutually exclusive scenarios: First, the hybrid zone may have moved and left the *M. m. musculus* Y-chromosome in the *M. m. domesticus* territory, or second, the *M. m. musculus* Y-chromosome might have escaped from a static hybrid zone

centre comprising autosomal and X-chromosomal markers and advanced up to 22km into the *M. m. domesticus* territory (Macholan *et al.*, 2008). It was proposed that a genetic conflict between the sexes may act together with a distortion of the sex-ratio to promote unidirectional sex-specific gene flow between the subspecies (Macholan *et al.*, 2008). The variability in marker cline patterns stresses the importance of replicated analyses for hybrid zone analyses. In the present study, we collected information on multiple hybrid zones to assess the properties of sex-specific marker clines in a speciation-prone European small mammal.

Voles in the genus *Microtus* have been undergoing the fastest radiation known in mammals and diversified into at least 65 species across the world within less than 2 million years (Fink *et al.*, 2010; Jaarola *et al.*, 2004). Despite very little morphological differentiation, karyotypic variation and differentiation into deep genetic lineages is relatively common within recognized species, suggesting the presence of cryptic species or ongoing speciation processes (Heckel *et al.*, 2005; Hellborg *et al.*, 2005; Jaarola *et al.*, 2004). In the common vole (*Microtus arvalis*), intraspecific divergence has resulted in four main, phenotypically cryptic evolutionary lineages in Europe (Western, Central, Italian and Eastern lineages; Heckel *et al.*, 2005). Divergence between these lineages based on population samples of mtDNA sequences and autosomal information was estimated to have coincided with or predated the last glacial maximum depending on the lineage resulting e.g. in more than 50,000 generations of divergence between the Western and Central lineages (Heckel *et al.*, 2005). An initial analysis of phylogeographic patterns in the region of secondary contact between lineages in the Alps demonstrated clear-cut parapatry of mtDNA lineages (Braaker and Heckel, 2009). Multi-locus nuclear microsatellite markers allow the allocation of individuals to genetic clusters whose

distributions are highly consistent with mtDNA lineages - except for first evidence of cytonuclear discordance in a few individuals close to the potential contact zones (Braaker and Heckel, 2009). The detected patterns are consistent with male-biased dispersal in the species (Hahne *et al.*, 2011; Schweizer *et al.*, 2007) and relatively complex processes of genetic erosion after colonization of the region (Braaker and Heckel, 2009). However, they may also be explained by an ongoing speciation process and the effect of sex-specific gene flow associated with yet undocumented partial reproductive isolation between the highly divergent evolutionary lineages in *M. arvalis*. With male-biased dispersal in the species, a selectively neutral secondary contact would show broader paternal marker clines than maternal ones. In a case involving selection e.g. a form of Haldane's rule, paternal marker clines would not be expected to be broader than maternal ones because they are transmitted by the heterogametic sex and thus gene flow would be expected to be restricted.

Here we fine-map the geographic location of the secondary contact zones between all three evolutionary lineages (Western, Central, Italian) of *M. arvalis* occurring in the region of the central Alps. We performed cline analyses of Y-chromosomal and mtDNA markers across these contact zones to quantify the relative contribution of sex-specific processes and the importance of non-neutral evolution linked to potential incipient speciation in the probably most abundant wild European mammal.

Materials and Methods

Sampling of transects

We investigated the different secondary contact zones in the region of the Alps between three evolutionary lineages of the common vole with a total number of 706 voles in transects of 50 to 160 km length consisting of 23 to 40 localities each (Figure 1, Table 2, Supplementary Information Table 1). The first transect (Jura) was sampled perpendicular to the Jura mountain ridge, where the Western and the Central lineages meet. The second transect (Grisons) was sampled along the Rhine valley, a contact zone of the Central and the Italian lineages. The third transect (Valais) comprises transitions from the Central to the Western and then to the Italian lineage along the Rhone valley. Samples were obtained as described in Braaker and Heckel (2009).

Molecular markers, RFLP analysis and sequencing

We first determined a diagnostic Y-chromosome marker for the Italian, Western and Central lineages as follows: 745 bp of intron 11 of the *SMCY* gene were sequenced (see below for amplification and sequencing protocol) in 71 voles from 31 populations across Europe, covering the range of the Western (Spain, France, United Kingdom and Belgium), Central (Switzerland, Germany, and Austria) and Italian (Italy, Switzerland) lineages. Six haplotypes were found across Europe with a phylogeographic structure mirroring the parapatric distribution of mitochondrial lineages in *M. arvalis* (Meier *et al.*, in prep.; Heckel *et al.*, 2005). These were termed Western Y (three haplotypes), Central Y (two haplotypes) and Italian Y (one haplotype) analogous to the mitochondrial lineages and genetic clusters in multi-locus nuclear

147 markers (Braaker and Heckel, 2009; Heckel *et al*, 2005). Three of these lineage-diagnostic Y
148 haplotypes defined by four substitutions and one insertion/deletion were found in the
149 transects studied here (Table 1; 2).

150 *SMCY11* fragments were amplified in a reaction volume of 25µl using the primers
151 SMCY11-f (5'-CTG CCC TGY RCC ATG CAT-3') and SMCY11-r (5'-TCC ACC TGT TSM AGR ACA T-3')
152 from Hellborg and Ellegren (2003) developed for *M. agrestis*. For PCR amplification, an initial
153 step of denaturation at 95°C for 2 min was followed by 40 cycles of denaturation at 95°C for 30
154 sec, annealing at 49°C for 1 min and extension at 72°C for 90 sec. A final step at 72°C for 10 min
155 was added to complete primer extension. All PCR amplifications were performed in thermal
156 cyclers PCR System 9700 (GeneAmp, Applied Biosystems, Rotkreuz, Switzerland) or PTC-100TM
157 (MJ Research, Watertown, MA). For sequencing, PCR products were cleaned with a GenElute™
158 PCR clean-up kit (Sigma-Aldrich, St. Louis, USA) and dissolved in 40µl bi-distilled water. Cycle
159 sequencing reactions were carried out using the Terminator Ready Reaction Mix "Big Dye" 3.1
160 from Applied Biosystems and the PCR primers. The reactions (10µl) were performed with the
161 following conditions: denaturation at 95°C for 50 sec, 30 cycles of 96°C for 10 sec, 57°C for 10
162 sec and 60°C for 4.5 min. PCR products were cleaned by precipitation according to the Applied
163 Biosystems manual, and separated and detected on an Applied Biosystems Prism 3100 Genetic
164 Analyser.

165 The mitochondrial Cytochrome *b* (*cytb*) gene was amplified following Fink *et al*. (2004;
166 2006). The allocation of *cytb* amplification products to one of the three lineages present in the
167 study region was performed by enzyme digestion with AFLIII and BsrI and Restriction Fragment

Length Polymorphism (RFLP) analysis. Samples from the Jura transect were analysed with AFLIII which cuts the Central, Italian and Eastern lineage *cytb*, but not the Western lineage. Samples from the Grisons transect were analysed with BsrI, an enzyme cutting *cytb* fragments of the Western, Central and Eastern lineage twice, whereas Italian samples are only cut once. The samples from the Valais transect were analysed subsequently with both enzymes to enable allocation to one of the three lineages present. As a control, 20 – 40 samples per transect were additionally sequenced (see Braaker and Heckel, 2009) to verify lineage allocation by RFLP. No disagreements were found. For completeness, published *cytb* data of individuals from the study region (Braaker and Heckel 2009; Fink *et al.*, 2004; Heckel *et al.*, 2005) were integrated into our analyses. Sequences were aligned using the Clustal W algorithm (Thompson *et al.*, 1997) implemented in BioEdit 7.0.5 (Hall, 1999) and revised manually. Database accession numbers will be included upon acceptance.

Cline analyses

Contact zones can be described by a sigmoid cline in allele frequencies (*p*) with a function of cline centre (*c*) and width (*w*; defined as the inverse of the maximum slope) following

$$p = (1 + \tanh [2(x-c)/w])/2,$$

with *x* equal to the position of the population along the cline (Szymura and Barton, 1986). Cline fitting analysis was performed by reducing the location data to one-dimensional transects. The frequency of a lineage in a population was plotted against distance (km) along the transects from the north-westernmost (Jura), the northernmost (Grisons) or the westernmost (Valais) locality. Tanh curves were then fitted for the *cytb* and *SMCY11* data separately using the 'Fit 1D

cline' operation in the program ANALYSE version 1.3 (Barton and Baird, 2002). Model parameters (centre and width) were computed using 2000 iterations from 20 different starting points. Likelihood profiles and two log-likelihood support limits (corresponding to 95% confidence intervals; Edwards, 1972) were explored using the 'cross-section' option. Cline coincidence (centres at the same position) and concordance (equal cline widths) were explored with a Likelihood Ratio Test (LRT) following the procedure used by Phillips *et al.* (2004) and adapted by Leache and Cole (2007).

Results

Lineage assignment and marker mismatches

Dedicated sampling along transects allowed the precise localisation of the intraspecific contact zones in *M. arvalis* and a general refinement of the distribution of mtDNA lineages compared to a previous study (Braaker and Heckel 2009). The distribution of Y-chromosomal lineages was in general agreement with the mtDNA contact zones but with highly significant differences in each transect (Figure 2; Table 3).

In the Jura transect, 34 of 126 males showed a mismatch between the maternal and the paternal marker of which all but one had a Central Y chromosome and Western mtDNA. In the Grisons transect, a mismatch between the maternal and paternal marker was detected in 41 of 104 males, all with Italian mtDNA and a Central Y chromosome. In the Valais transect, our analyses showed a double transition between lineages along the Rhone valley for both markers. Despite extensive sampling attempts, no samples or evidence of the presence of *M.*

arvalis could be obtained between kilometres 18 and 75 of the transect separating the Central and Western mtDNA (Figure 1). Suitable habitats for the common vole are very scarce in this region due to intensive human usage. A mismatch between the maternal and paternal marker was detected in 28 out of 55 males sampled along the Valais transect. Two males had Western mtDNA and a Central Y chromosome while 26 had Western mtDNA and an Italian Y chromosome. In the Valais transect, the Western Y chromosome was found exclusively in combination with Western mtDNA.

Cline analyses

Cline shape analyses revealed significant cline centre shifts (non-coincidence) in all three transects and differences in cline widths (non-concordance) in two of the three transects (Figure 2; Table 3). In the Jura transect, the LRT revealed a significant north-westward shift of the Y-chromosomal cline relative to the mtDNA cline ($\chi^2_1 = 129.398$; for $df=1$ the critical value at $p = 0.01$ is 6.635). Moreover, the mtDNA cline was significantly wider ($\chi^2_1 = 34.342$). The Grisons transect also displayed non-coincident ($\chi^2_1 = 69.990$) and non-concordant ($\chi^2_1 = 56.774$) clines. The Y-chromosomal cline was narrower than the mtDNA cline and the centre shifted to the South. In the Valais transect, analyses were performed separately for the two transition zones (Central-Western and Western-Italian). Centres and widths of the paternal and maternal cline of the Central-Western contact zone did not differ significantly in this part of the transect containing mostly unsuitable vole habitats. However, several males with a Central Y chromosome were detected higher up in the Rhone valley at kilometre 78 in a population with only Western mtDNA which was several dozens of kilometres away from the nearest known

Central Y males along the transect (Figure 2). Coincidence of the Western-Italian contact zone was highly significantly rejected ($\chi^2_1 = 52.890$), whereas there was no evidence that the widths of the clines differed ($\chi^2_1 = 0.000$). Further analyses according to age classes of voles provided no evidence of differences in the position of centres or widths of clines between adults and juveniles in any transect (details not shown).

Discussion

Our analyses of sex-specific genetic markers across multiple contact zones revealed strong evidence for partial reproductive isolation between highly divergent evolutionary lineages within the common vole, *M. arvalis*. Cline analyses strongly suggest sex-specific processes acting in the contact zones consistent with the effect of a unidirectional manifestation of Haldane's rule.

Evidence for selection

Very narrow clines for the Y chromosome in comparison to mtDNA suggest the action of selection on common vole males. The width of a cline across a hybrid zone is theoretically shaped by dispersal which tends to widen it and selection which may narrow it (Barton and Hewitt, 1985). In *M. arvalis*, dispersal is male-biased like in most mammals (Hahne *et al.*, 2011; Hamilton *et al.*, 2005; Schweizer *et al.*, 2007). Thus without the action of selection paternally inherited markers are expected to show wider clines across the contact zones than maternally inherited ones. At odds with this neutral expectation, Y chromosome clines were significantly narrower or at most equal to mtDNA clines, suggesting selective forces acting on males. At

present, the mechanisms leading to this pattern are unknown (see below) but it is compatible with Haldane's rule predicting negative fitness effects for the heterogametic sex due to recessive X-linked mutations or epistatic interactions between the Y chromosome and another part of the genome (Coyne and Orr, 1998; 2004; Haldane, 1922; Turelli and Orr, 2000).

These patterns in *M. arvalis* are similar to other small mammal systems where speciation is ongoing or was recently completed but where evolutionary divergence is much older and phenotypic differentiation between the taxa has been established. For example, in the European hybrid zone between *Mus m. musculus* and *Mus m. domesticus* (diverged >700 000 generations ago; She *et al.*, 1990) the transition of sex chromosomes is similar in width to *M. arvalis* (e.g. < 4km) and mtDNA clines are also wider (e.g. Dod *et al.*, 1993; Macholan *et al.*, 2007; 2008; Raufaste *et al.*, 2005). This is likely a consequence of Haldane's rule in place in the hybrid zone which may – among other factors - prevent the mouse Y chromosome cline width to increase (Britton-Davidian *et al.*, 2005; but see Turner *et al.*, 2012). The rapidly speciating *Microtus* genus harbours several other examples of restricted hybridization and evidence for selection acting prevalently on males - typically associated with morphological and karyotypic differences between the involved sibling species or subspecies (Beysard *et al.*, 2012; Bulatova *et al.*, 2010; Jaarola *et al.*, 1997; Meier *et al.*, 1996). Among the cryptic evolutionary lineages in *M. arvalis* however, partial reproductive isolation appears to be additionally acting in an unusual unidirectional manner.

Asymmetrical sex-specific processes

The second sex-specific process across the contact zones is the shift between maternal and paternal clines leading to an asymmetrical pattern of mismatches. These mismatches may be the result of two main processes: introgression of mtDNA and/or introgression of the Y chromosome. First, a range expansion of one lineage may have left behind a trail of introgressed mtDNA in the other lineage (see Buggs, 2007 for a review). Simulation studies have shown that a replacement of one taxon by another may result in the introgression of selectively neutral local genes into the invading taxon and that markers transmitted by the least dispersing sex would be most introgressed (Currat *et al.*, 2008; Petit and Excoffier, 2009). In our case, this scenario would have resulted in asymmetrical introgression of mtDNA. However, this scenario would require a general fitness advantage of one lineage over the other, for which there is currently no support given the apparent absence of phenotypic differences between the lineages. mtDNA might also introgress due to a selective advantage into other taxa (see examples in Currat *et al.*, 2008 and Petit and Excoffier, 2009) but there is no evidence of mtDNA introgression or of a selective advantage of particular mtDNA lineages (see Braaker and Heckel, 2009; Fink *et al.*, 2004; Fischer *et al.*, 2011; Hamilton *et al.*, 2005). Additionally, very high diversity of mtDNA haplotypes in populations and lineages (see Braaker and Heckel, 2009; Heckel *et al.*, 2005) argues against a recent mtDNA sweep, but additional analyses together with the nuclear genome will be necessary for dedicated testing of mtDNA introgression in the hybrid zones.

The alternative main explanation for asymmetrical mismatches is the potential existence of asymmetric pre- or post-zygotic mechanisms leading to an introgression of the Y chromosome in the opposite direction. Pre-zygotic mechanisms might consist e.g. in mating

297 preferences of females from both lineages for a certain Y-chromosomal lineage or in specific
298 male-male interactions (e.g. aggressive behaviour; Roubertoux *et al.*, 1994). A potential post-
299 zygotic mechanism in *M. arvalis* would be a unidirectional manifestation of Haldane's rule with
300 incompatibilities occurring only in backcrosses involving hybrid males with one sex
301 chromosome type but not the other. Mating and breeding experiments involving these *M.*
302 *arvalis* lineages will allow to test for behavioural mechanisms or genetic incompatibilities acting
303 as selective forces.

304 A possible unidirectional manifestation of Haldane's rule was already suggested by
305 Gavrillets (1997) based on theoretical work. Non-coincident marker clines similar to the ones in
306 *M. arvalis* in a particular region of the European house mouse hybrid zone are compatible with
307 such a process as well (Macholan *et al.*, 2008). The *Mus m. musculus* Y chromosome and to a
308 lesser extent X chromosomal markers and mtDNA are here more introgressed into the *Mus m.*
309 *domesticus* territory than the rest of the genome (Macholan *et al.*, 2008; 2011). The authors
310 suggested that this unusual pattern resulted from genetic conflict between the sexes which
311 may promote unidirectional sex-specific gene flow between subspecies. To date, this pattern
312 has only been detected in one of the studied transects and the general importance of stochastic
313 vs. deterministic effects in the house mouse hybrid zone deserves further investigation (see
314 Macholan *et al.*, 2011; Turner *et al.*, 2012). A potentially promising similar case was found in
315 Scandinavia, where 15 mice from seven *M. m. domesticus* populations were introgressed with
316 the *M. m. musculus* Y-chromosome (Jones *et al.*, 2010). Our analyses, however, display a highly
317 consistent pattern of very sharp Y-chromosomal clines shifted relative to mtDNA clines in
318 several hybrid zones between the morphologically cryptic lineages of *M. arvalis*. At present, it is

unclear if similar forces are acting in the different contact zones but additional X-chromosomal and autosomal markers will allow a more detailed characterisation of introgression patterns in the future. This may also reveal particular differences between hybrid zones, potentially depending on the evolutionary lineages involved and their level of divergence.

Conclusion

Our analyses of multiple hybrid zones provide evidence for sex-specific processes which are compatible with strong selection acting at the contact between evolutionary lineages in the common vole. The mechanisms contributing to partial reproductive isolation and the extent of actual hybridization in natural populations remain currently unknown. However, more detailed population genetic analyses covering the nuclear genome widely combined with classical experimental crosses and mate-choice experiments and the analysis of less-diverged lineages (e.g. Eastern) have the potential to inform us about causes and consequences of asymmetric gene flow in hybrid zones of the probably most abundant wild European mammal.

Acknowledgements

We are grateful to Laurent Excoffier and the University of Bern for ongoing support of the vole projects. We thank the editor and three reviewers for helpful suggestions, Irene Keller and Mathieu Foll for comments on the manuscript, Susanne Tellenbach for technical assistance, Joana Meier for unpublished data on Y-chromosomal variability, and Sonja Braaker, Martin

Fischer and Birgit Thorenz for kindly providing access to specimens. This study was supported in part by grants from the Berne University Research Foundation and by the Swiss National Science Foundation number 31003A-127377 to GH.

Conflict of interest

The authors declare no conflict of interest.

References

- Baird SJE, Ribas A, Macholán M, Albrecht T, Piálek J, Goüy de Bellocq J (2012). Where are the wormy mice? A reexamination of hybrid parasitism in the European House mouse hybrid zone. *Evolution* **66**: 2757-2772.
- Barton NH, Baird SJE (2002). ANALYSE, Version 1.3: Freeware: available from <http://helios.bto.edsac.uk/evolgen/Mac/Analyse>.
- Barton NH, Hewitt GM (1985). Analysis of hybrid zones. *Annu Rev Ecol Syst* **16**: 113-148.
- Barton NH, Hewitt GM (1989). Adaptation, speciation and hybrid zones. *Nature* **341**: 497-503.
- Beysard M, Perrin N, Jaarola M, Heckel G, Vogel P (2012). Asymmetric and differential gene introgression at a contact zone between two highly divergent lineages of field voles (*Microtus agrestis*). *J Evol Biol* **25**: 400-408.
- Braaker S, Heckel G (2009). Transalpine colonisation and partial phylogeographic erosion by dispersal in the common vole (*Microtus arvalis*). *Mol Ecol* **18**: 2518-2531.

359 Britton-Davidian J, Fel-Clair F, Lopez J, Alibert P, Boursot P (2005). Postzygotic isolation between
360 the two European subspecies of the house mouse: estimates from fertility patterns in
361 wild and laboratory-bred hybrids. *Biol J Linn Soc* **84**: 379-393.

362 Buggs RJA (2007). Empirical study of hybrid zone movement. *Heredity* **99**: 301-312.

363 Bulatova NS, Potapov SG, Lavrenchenko LA (2010). Genomic versus chromosomal polytypy in
364 studies of mitochondrial and nuclear DNA markers in the *Microtus arvalis* group. *Russ J*
365 *Genet* **46**: 586-594.

366 Coyne JA, Orr HA (1998). The evolutionary genetics of speciation. *Phil Trans R Soc B Biol Sci* **353**:
367 287-305.

368 Coyne JA, Orr HA (2004). *Speciation*. Sinauer: Sunderland.

369 Currat M, Ruedi M, Petit RJ, Excoffier L (2008). The hidden side of invasions: Massive
370 introgression by local genes. *Evolution* **62**: 1908-1920.

371 Dod B, Jermini LS, Boursot P, Chapman VH, Nielsen JT, Bonhomme F (1993). Counterselection
372 on sex-chromosomes in the *Mus musculus* European hybrid zone. *J Evol Biol* **6**: 529-546.

373 Edwards AWF (1972). *Likelihood*. Cambridge University Press: Cambridge, U. K.

374 Fink S, Excoffier L, Heckel G (2004). Mitochondrial gene diversity in the common vole *Microtus*
375 *arvalis* shaped by historical divergence and local adaptations. *Mol Ecol* **13**: 3501-3514.

376 Fink S, Excoffier L, Heckel G (2006). Mammalian monogamy is not controlled by a single gene.
377 *PNAS* **103**: 10956-10960.

378 Fink S, Fischer MC, Excoffier L, Heckel G (2010). Genomic scans support repetitive continental
379 colonization events during the rapid radiation of voles (Rodentia: *Microtus*): the utility of
380 AFLPs versus mitochondrial and nuclear sequence markers. *Syst Biol* **59**: 548-572.

381 Fischer MC, Foll M, Excoffier L, Heckel G (2011). Enhanced AFLP genome scans detect local
382 adaptation in high-altitude populations of a small rodent (*Microtus arvalis*). *Mol Ecol* **20**:
383 1450-1462.

384 Gavrillets S (1997). Hybrid zones with Dobzhansky-type epistatic selection. *Evolution* **51**: 1027-
385 1035.

386 Ganem G, Lital C, Lenormand T (2008). Variation in mate preference across a house mouse
387 hybrid zone. *Heredity* **100**: 594-601.

388 Good JM, Handel MA, Nachman MW (2008). Asymmetry and polymorphism of hybrid male
389 sterility during the early stages of speciation in house mice. *Evolution* **62**: 50-65.

390 Hahne J, Jenkins T, Halle S, Heckel G (2011). Establishment success and resulting fitness
391 consequences for vole dispersers. *Oikos* **120**: 95-105.

392 Haldane J (1922). Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics* **12**: 101-
393 109.

394 Hall TA (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis
395 program for Windows 95/98/NT. *Nuc Acids Symp Ser* **41**: 95-98.

396 Hamilton G, Currat M, Ray N, Heckel G, Beaumont M, Excoffier L (2005). Bayesian estimation of
397 recent migration rates after a spatial expansion. *Genetics* **170**: 409-417.

398 Heckel G, Burri R, Fink S, Desmet JF, Excoffier L (2005). Genetic structure and colonization
399 processes in European populations of the common vole, *Microtus arvalis*. *Evolution* **59**:
400 2231-2242.

401 Hellborg L, Ellegren H (2003). Y chromosome conserved anchored tagged sequences (YCATS) for
402 the analysis of mammalian male-specific DNA. *Mol Ecol* **12**: 283-291.

403 Hellborg L, Gunduz I, Jaarola M (2005). Analysis of sex-linked sequences supports a new
404 mammal species in Europe. *Mol Ecol* **14**: 2025-2031.

405 Jaarola M, Martinkova N, Gunduz I, Brunhoff C, Zima J, Nadachowski A *et al.* (2004). Molecular
406 phylogeny of the speciose vole genus *Microtus* (*Arvicolinae*, *Rodentia*) inferred from
407 mitochondrial DNA sequences. *Mol Phylogenet Evol* **33**: 647-663.

408 Jaarola M, Tegelström H, Fredga K (1997). A contact zone with noncoincident clines for sex-
409 specific markers in the field vole (*Microtus agrestis*). *Evolution* **51**: 241-249.

410 Jones EP, van der Kooij J, Solheim R, Searle JB (2010). Norwegian house mice (*Mus musculus*
411 *musculus/domesticus*): distributions, routes of colonization and patterns of hybridization. *Mol Ecol*
412 **19**: 5252-5264.

413 Leache AD, Cole CJ (2007). Hybridization between multiple fence lizard lineages in an ecotone:
414 locally discordant variation in mitochondrial DNA, chromosomes, and morphology. *Mol*
415 *Ecol* **16**: 1035-1054.

416 Macholan M, Baird SJE, Dufkova P, Munclinger P, Bimova BV, Pialek J (2011). Assessing
417 multilocus introgression patterns: a case study on the mouse X chromosome in Central
418 Europe. *Evolution* **65**: 1428-1446.

419 Macholan M, Baird SJE, Munclinger P, Dufkova P, Bimova B, Pialek J (2008). Genetic conflict
420 outweighs heterogametic incompatibility in the mouse hybrid zone? *BMC Evol Biol* **8**: e14.

421 Macholan M, Munclinger P, Sugerkova M, Dufkova P, Bimova B, Bozikova E *et al.* (2007).
422 Genetic analysis of autosomal and X-linked markers across a mouse hybrid zone.
423 *Evolution* **61**: 746-771.

424 Meier MN, Golenishchev FN, Radzhabli SI, Sablina OL (1996). Gray voles (subgenus *Microtus*) of
425 the fauna of Russia and adjacent territories. *Trudy Zoolog ichesкого insituta (Collection of*
426 *Scientific Papers of Zoological Institute) St Petersburg: Ross Akad Nauk.*

427 Mihola O, Trachtulec Z, Vlcek C, Schimenti JC, Forejt J (2009). A mouse speciation gene encodes
428 a meiotic histone H3 methyltransferase. *Science* **323**: 373-375.

429 Petit E, Balloux F, Excoffier L (2002). Mammalian population genetics: why not Y? *Trends Ecol*
430 *Evol* **17**: 28-33.

431 Petit RJ, Excoffier L (2009). Gene flow and species delimitation. *Trends Ecol Evol* **24**: 386-393.

432 Phillips BL, Baird SJE, Moritz C (2004). When vicars meet: A narrow contact zone between
433 morphologically cryptic phylogeographic lineages of the rainforest skink, *Carlia*
434 *rubrigularis*. *Evolution* **58**: 1536-1548.

435 Raufaste N, Orth A, Belkhir K, Senet D, Smadja C, Baird SJE *et al.* (2005). Inferences of selection
436 and migration in the Danish house mouse hybrid zone. *Biol J Linn Soc* **84**: 593-616.

437 Roubertoux PL, Carlier M, Degrelle H, Haasdupertuis MC, Phillips J, Moutier R (1994). Co-
438 segregation of intermale aggression with the pseudoautosomal region of the Y-
439 chromosome in mice. *Genetics* **136**: 225-230.

440 Schilthuizen M, Giesbers MCWG, Beukeboom LW (2011). Haldane's rule in the 21st century.
441 *Heredity* **107**: 95-102.

442 Schweizer M, Excoffier L, Heckel G (2007). Fine-scale genetic structure and dispersal patterns in
443 the common vole (*Microtus arvalis*). *Mol Ecol* **16**: 2463-2473.

She JX, Bonhomme F, Boursot P, Thaler L, Catzeflis F (1990). Molecular phylogenies in the genus *Mus* - comparative analysis of electrophoretic, scnDNA hybridization, and mtDNA RFLP data. *Biol J Linn Soc* **41**: 83-103.

Szymura JM, Barton NH (1986). Genetic-analysis of a hybrid zone between the fire-bellied toads, *Bombina bombina* and *Bombina variegata*, near Cracow in southern Poland. *Evolution* **40**: 1141-1159.

Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997). The CLUSTAL X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* **25**: 4876-4882.

Turelli M, Begun DJ (1997). Haldane's rule and X-chromosome size in *Drosophila*. *Genetics* **147**: 1799-1815.

Turelli M, Orr HA (2000). Dominance, epistasis and the genetics of postzygotic isolation. *Genetics* **154**: 1663-1679.

Turner LM, Schwahn DJ, Harr B (2012). Reduced male fertility is common but highly variable in form and severity in a natural house mouse hybrid zone. *Evolution* **66**: 443-458.

Yannic G, Basset P, Hausser J (2008). A hybrid zone with coincident clines for autosomal and sex-specific markers in the *Sorex araneus* group. *J Evol Biol* **21**: 658-667.

Data accessibility:

DNA sequences will be deposited in Genbank and accession numbers will be provided upon acceptance.

Table captions:

Table 1:

Haplotypes and polymorphic positions in the Y-chromosomal marker (intron SMCY 11; 745 bp) in the studied transects across contact zones in *M. arvalis*. The position of the substitutions and insertion/deletion are indicated in the first row. The number of individuals sequenced per haplotype is displayed under N.

Table 2:

Summary of the three transects across contacts between evolutionary lineages in *Microtus arvalis*. Positions of transects and sampling localities are shown in Figure 1. The Valais transect along the Rhone valley contains a double transition from the Central (C) through the Western (W) to the Italian (It) lineage. Sample sizes are given for mtDNA (*cytb*) and Y-chromosomal sequences.

Table 3:

Contact zone cline parameters and likelihood-ratio test results of the null hypothesis that cline centres and widths do not differ significantly between the maternal and parental marker for a given transect. Maximum likelihood estimates of the position of cline centres and their widths are given in kilometres and confidence intervals (two log-likelihood units \approx 95% c.i.) are shown in parentheses. $\ln L_u$ is the unconstrained log-likelihood support. $\ln L_c$ is the maximum constrained likelihood in an analysis where cline centre or width was held constant at intervals while allowing the other parameter (width or centre) to vary freely. An asterisk indicates

488 rejection of the null hypothesis at $\alpha = 0.01$ (critical χ^2 value = 6.635; one degree of freedom).

489 Estimates for centres and widths in the Valais transect are given separately for the two

490 different transition zones (Central-Western and Western-Italian).

491

492 Table 1:

Haplotype	N	135	179	298	339	608
Western Y1	42	C	T	C	G	T
Italian Y	51	T	A	C	G	T
Central Y1	192	C	T	T	-	C

493

494 Table 2:

Transect	Lineage transition	First/last locality	Length [km]	Localities	N (<i>Cyt b</i>)	N (<i>SMCY11</i>)
Jura	Western-Central	Cerniébaud (F)/ Les Cullayes	51	40	281 (178 W - 103 C)	126 (36 W - 90 C)
Grisons	Central-Italian	Diepoldsau/ Jochalp	71	26	287 (133 C - 154 It)	104 (90 C - 14 It)
Valais	Central - Western Western - Italian	Aigle/ Villa	159	23	138 (40 C - 79 W - 19 It)	55 (12 C - 6 W - 37 It)

496 Table 3:

	Parameter	cytb	SMCY11	$\sum \ln L_u$	$\sum \ln L_c$	$\Delta = \sum \ln L_c - \sum \ln L_u$	$\chi^2 = 2 \Delta$
Jura	Cline centre	28.98 (27.68 - 30.38)	17.5 (17.15-18.05)	-63.7	-128.399	64.699	*129.398
	Cline width	16.81 (13.31 - 21.91)	1.59 (0.89-3.34)	-63.7	-80.871	17.171	*34.342
	$\ln L_u$	-53.973	-9.727				
Grisons	Cline centre	49.32 (46.12-52.22)	65.47 (64.47-67.37)	-76.6	-111.595	34.995	*69.990
	Cline width	39.47 (31.00-50.97)	0.25 (0.0-2.0)	-76.6	-104.987	28.387	*56.774
	$\ln L_u$	-76.6	-0.000				
Valais (C-W)	Cline centre	28.4 - 70 (17.7-75.7)	74.23 (45.8-77.9)	-0.701	-0.702	0.001	0.002
	Cline width	2.1 - 10.8 (0.01-37.7)	10.19 (2.2-56.6)	-0.701	-0.701	0.000	0.000
	$\ln L_u$	-0.000	-0.701				
Valais (W-I)	Cline centre	100.3 (99.9-105.2)	78.4 (78.1-81.6)	-0.001	-26.446	26.445	*52.890
	Cline width	0.1 - 2 (0.01-10.9)	0.1 - 1.1 (0.01-9.2)	-0.001	-0.001	0.000	0.000
	$\ln L_u$	-0.001	-0.000				

1 **Titles and legends to figures:**

2

3 **Figure 1:**

4 Map of Switzerland showing the sampling localities of *Microtus arvalis* samples
5 included in cline analyses of the three transects. Transects are described in Table 1.
6 Localities are designated according to the *cytb* lineage they harbour (white = Central,
7 grey = Western, black = Italian). The Valais transect along the Rhone valley contains a
8 double transition from the Central through the Western to the Italian lineage. Mixed
9 populations (see Figure 2) are indicated by bi-coloured circles. Localities closer than
10 two kilometres to each other were merged for this figure.

11

12 **Figure 2:**

13 Plots of lineage frequencies in populations versus position (km) along the transects
14 Jura, Grisons and Valais for mtDNA (white circles) and Y-chromosomal (grey
15 triangles) markers. Sizes of symbols are proportional to the log number of samples.
16 Bold lines are fitted clines for mtDNA and dashed lines for Y-chromosomal markers.
17 Despite a sampling gap in the Valais transect, an admixed population was detected
18 at km 78, pointing to at least partial contact between the Western and the Central
19 lineage in this transect. See Table 2 for cline properties and statistical comparisons.

Fig.1




Fig. 2

